



Product Information Sheet

H437 Hosta Rooting Medium Stage III Medium

Properties

Form:	Powder
Appearance:	White to Yellow
Application:	Plant Tissue Culture
Solubility:	Partially Soluble in Cold Water. Fully Soluble in Hot/Boiling Water.
Typical Working Concentration:	43.23 g/L
Storage Temp:	2-6°C
Storage Temp of Stock Solution:	Preparation of concentrated solutions is not recommended as insoluble precipitates may form.
Other Notes:	Contains the macro- and micronutrients as described by Murashige and Skoog (1962) and the vitamins described by Linsmaier and Skoog (1965). Also contains agar and sucrose. Plant Tissue Culture Tested.

Formula (mg/L)

Ammonium Nitrate	1650
Boric Acid	6.2
Calcium Chloride, Anhydrous	332.2
Cobalt Chloride•6H ₂ O	0.025
Cupric Sulfate•5H ₂ O	0.025
Disodium EDTA •2H ₂ O	37.26
Ferrous Sulfate•7H ₂ O	27.8
Magnesium Sulfate, Anhydrous	180.7
Manganese Sulfate•H ₂ O	16.9
Molybdic Acid (Sodium Salt)• 2H ₂ O	0.25
Potassium Iodide	0.83

Potassium Nitrate	1900
Potassium Phosphate, Monobasic	300
Sodium Phosphate, Monobasic	170
Zinc Sulfate•7H ₂ O	8.6
Agar	8000
6-Benzylaminopurine (BA)	0.1
Casein, Enzymatic Hydrolysate	500
myo-Inositol	100
α-Naphthaleneacetic Acid	0.5
Sucrose	30,000
Thiamine•HCl	0.4

Application Notes

Plant species: Hosta

References

- Linsmaier, EM and F Skoog. 1965. Organic growth factor requirements of tobacco tissue cultures. *Physiol. Plant.* 18: 100-127.
- Murashige, T and F Skoog. 1962. A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Physiologia Plantarum.* 15: 473-497.

Revised 4/2013