



Product Information Sheet

K4751 Kanamycin Solution (100 mg/mL)

Synonyms: O-3-Amino-3-deoxy- α -D-glucopyranosyl-(1 \rightarrow 6)-O-[6-amino-6-deoxy- α -D-glucopyranosyl-(1 \rightarrow 4)]-2-deoxy-D-Streptamine Monosulfate

CAS: 25389-94-0

Formula: $C_{18}H_{36}N_4O_{11} \cdot H_2SO_4$

Mol. Weight: 582.6

Properties

Form: Liquid


Appearance: Colorless, Clear Liquid

Application: Plant Tissue Culture Antibiotic

Solubility: Miscible with Water

Storage Temp: 2 to 6 °C

Other Notes: Plant Tissue Culture Tested



Application Notes

Kanamycin is an aminoglycoside antibiotic derived from *Streptomyces kanamyceticus*. Kanamycin has similar mode of action to those of gentamicin in which it inhibits protein synthesis and elicits miscoding by binding to the 30S subunit and sometimes the 50S subunit of the bacterial ribosome. Kanamycin is effective against Gram-positive and Gram-negative bacteria.^{2,3} It has been reported that cross-resistance occurs between kanamycin and neomycin, framycetin, and paromomycin, and partial cross-resistance between kanamycin and streptomycin.²

It has been reported that the minimum inhibitory concentration (MIC) of kanamycin for *Staphylococcus aureus* is 3.5 mg/L and *Escherichia coli* is 4.5 mg/L.⁴

PhytoTechnology Laboratories® also carries kanamycin solution at 50 mg/mL (Product No. K586) and powder form of kanamycin monosulfate (Product No. K378).

Please Note: While PhytoTechnology Laboratories® tests each lot of this product with two or more plant cell/ tissue culture lines, it is the sole responsibility of the purchaser to determine the appropriateness of this product for the specific plants that are being cultured and applications that are being used.

References

1. Merck **13**, 5299
2. *Martindale: The Complete Drug Reference*, 35th ed., Paul S. Blake, Ed. (Royal Pharmaceutical Society, 2007), p. 260.
3. Divyashree BC, Vishwanatha T, Spoorthi N Jain, Reena V, Aishwarya Sampath, Siddhalingeswara KG, Venugopal N. 2012. Evaluation of in vitro synergy between ampicillin and kanamycin against *Staphylococcus aureus*. *Journal of Drug Delivery & Therapeutics*. Vol 2(4). Pp. 144-146.
4. Pereira, Ana Margarida, Ana Cristina Abreu, and Manuel Simoes. 2012. Action of kanamycin against single and dual species biofilms of *Escherichia coli* and *Staphylococcus aureus*. *Journal of Microbiology Research*. Vol 2(4). Pp. 84-88.

PhytoTechnology Laboratories®

P.O. Box 12205; Shawnee Mission, KS 66282-2205

Phone: 1-888-749-8682 or 1-913-341-5343; Fax: 1-888-449-8682 or 1-913-341-5442

Web Site: www.phytotechlab.com

© 2014 PhytoTechnology Laboratories®