




Product Information Sheet

M569
MUG



Synonym: 4-Methylumbelliferyl- β -D-glucuronide Trihydrate

CAS: 6160-80-1

Formula: $C_{16}H_{16}O_9 \cdot 3H_2O$

Molecular Wt: 406.4

Properties

Form: Powder

Appearance: White Fluffy Powder

Application: Molecular Biology

Solubility: DMSO

Typical Working

Concentration:

Storage Temp: -20 to 0 °C

Storage Temp of

Stock Solution: -20 to 0 °C

Application Notes

Fluorescent substrate for β -galactosidase.

References

Bai F and DeMason DA (2008) Hormone interactions and regulation of PsPK2::GUS compared with DR5::GUS and PID::GUS in *Arabidopsis thaliana*. *Am. J. Bot.* vol. 95 no. 2 133-145
doi:[10.3732/ajb.95.2.133](https://doi.org/10.3732/ajb.95.2.133)

Blázquez M (2007) Quantitative GUS Activity Assay of Plant Extracts. *Cold Spring Harb Protoc*; 2007; doi:[10.1101/pdb.prot4690](https://doi.org/10.1101/pdb.prot4690)

Blázquez M (2007) Quantitative GUS Activity Assay in Intact Plant Tissue. *Cold Spring Harb Protoc*; doi:[10.1101/pdb.prot4688](https://doi.org/10.1101/pdb.prot4688)

Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*, **72**, 248-254.

PhytoTechnology Laboratories®

P.O. Box 12205 • Shawnee Mission, KS • 66282-2205

Phone: 1-913-341-5343 or 1-888-749-8682 (U.S. Only) Fax: 1-913-341-5442 or 1-888-449-8682 (U.S. Only)

Web Site: www.phytotechlab.com

© 2011 PhytoTechnology Laboratories®



Product Information Sheet

- Conner AJ, Mlynarova L, Stiekema WJ and Nap JP (1999) Gametophytic expression of GUS activity controlled by the potato Lhca3.St.1 promoter in tobacco pollen. *Journal of Experimental Botany*, Vol. 50, No. 338, pp. 1471–1479.
- Curi GC, Chan RL and Gonzalez, DH (2005) The leader intron of Arabidopsis thaliana genes encoding cytochrome c oxidase subunit 5c promotes high-level expression by increasing transcript abundance and translation efficiency. *J. Exp. Bot.* 56 (419): 2563-2571. doi: [10.1093/jxb/eri250](https://doi.org/10.1093/jxb/eri250)
- Fior S and Gerola PD (2009) Impact of ubiquitous inhibitors on the GUS gene reporter system: evidence from the model plants Arabidopsis, tobacco and rice and correction methods for quantitative assays of transgenic and endogenous GUS. *Plant Methods* 5:19. doi:[10.1186/1746-4811-5-19](https://doi.org/10.1186/1746-4811-5-19)
- Gallagher, S.R. (1992) Quantitation of GUS activity by fluorometry. In GUS Protocols: using the GUS gene as a reporter of gene expression (Gallagher, S.R., ed): Academic Press, Inc., New York, pp. 47-59.
- Herd S, Christensen MJ, Saunders K, Scott DB and Schmid J (1997) Quantitative assessment of in planta distribution of metabolic activity and gene expression of an endophytic fungus. *Microbiology* 143, 267-275
- Janssen, B.J., and R.C. Gardner, 1989. Localized transient expression of GUS in leaf discs following co-cultivation with Agrobacterium, *Plant Mol. Biol.* 14: 61-72.
- Jefferson, R.A., Kavanagh, T.A. and Bevan, M.W. (1987) GUS fusions: B-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.*, 6, 3901-3907.
- Jefferson, R.A. 1987. Assaying chimeric genes in plants, the GUS gene fusion system, *Plant Mol. Biol. Rep.* 5: 387-405.
- Pamphilel JA, da Rochal CL, Azevedoll JL (2004) Co-transformation of a tropical maize endophytic isolate of Fusarium verticillioides (synonym F. moniliforme) with gusA and nia genes. *Genet. Mol. Biol.* vol.27 no.2; Pp. 253-258. doi: [10.1590/S1415-47572004000200021](https://doi.org/10.1590/S1415-47572004000200021)
- Wenzl P, Wong L, Kwang-won K and Jefferson RA (2005) A Functional Screen Identifies Lateral Transfer of β -Glucuronidase (gus) from Bacteria to Fungi. *Mol Biol Evol* 22 (2): 308-316. doi:[10.1093/molbev/msi018](https://doi.org/10.1093/molbev/msi018)