

T839

## Terrestrial (Cypripedium) Orchid Medium w/ Agar & w/o Ammonium Nitrate 400 mg/L Calcium Nitrate + 400 mg/L Casein

## **Properties:**

Form: Powder

Appearance: White to Yellow powder

Application: Orchid Culture

Solubility: Partially Soluble in Water

Typical Working Concentration:

27.44 g/L

Storage Temp: 2-8°C

Storage Temp of Preparation of concentrated solutions is not recommended as insoluble precipitates may

Stock Solution: form

Other Notes: Contains a modification of the macro- and micronutrients, glucose, and agar as described

by Steele (1996)

Contains 400 mg/L Calcium Nitrate and 400 mg/L Casein

Without Ammonium Nitrate

pH = 5.0 - 6.0

Formula [mg/L]:

Ammonium Citrate	19
Boric Acid	0.5
Calcium Nitrate	400
Cupric Sulfate•5H <sub>2</sub> O	0.025
Ferric Ammonium Citrate	25
Magnesium Sulfate, Anhydrous	97.69
Manganese Sulfate•H <sub>2</sub> O	1.54
Molybdic Acid, Sodium Salt•2H <sub>2</sub> O	0.02

Potassium Chloride	100
Potassium Iodide	0.1
Potassium Nitrate	200
Potassium Phosphate, Monobasic	200
Zinc Sulfate•7H <sub>2</sub> O	0.5
Agar	6000
Casein, Enzymatic Hydrolysate	400
D-Glucose	20,000

## **Application Notes:**

Plant Tissue Culture Tested

Plant species: Cypripedium and other terrestrial orchid species

Developed for germination and growth *Cypripedium reginae* (Harvais 1982). Ammonium nitrate is omitted so that it can be supplemented at 500-1000 mg/L with other *Cypripedium* sp. (Steele 1996).

## References:

Harvais G (1982) An improved culture medium for growing the orchid Cypripedium reginae axenically. *Can. J. Bot.* 60(12):2547-2555

Steele, WK. (1996). Large Scale Seedling Production of North American Cypripedium Species. In: C. Allen, Editor, North American Native Terrestrial Orchids. Propagation and Production Conf. Proc., May 16 & 17, 1996. pp 11-26.